

On Local Symmetric Order Parameters of Vortex Lattice States

Masa-aki OZAKI*, Yoshiki HORI[†] and Akira GOTO[‡]

**Department of Physics, Kochi University, Kochi 780-8072, Japan*

^{†,‡}*Kochi National College of Technology, Nankoku 783-8505, Japan*

(February 1, 2008)

This paper gives a new refined definition of local symmetric order parameters (OPs) (s -wave, d -wave and p -wave order parameters) of vortex lattice states for singlet superconductivity. s -wave, d -wave and p -wave OPs at a site (m, n) are defined as A , B and E representations of the four fold rotation C_4 at the site (m, n) of nearest neighbor OPs $\langle a_{(m,n)\downarrow} a_{(m+1,n)\uparrow} \rangle$ etc. The new OPs have a well defined nature such that an OP (e.g. d -wave) at the site obtained under translation by a lattice vector (of the vortex lattice) from a site (m, n) is expressed by the corresponding OP (e.g. d -wave) at the site (m, n) times a phase factor. The winding numbers of s -wave and d -wave OPs are given.

Growing interest has been focused on microscopic works on the basis of the Bogoliubov de Gennes (BG) equation of an isolated vortex¹ or a vortex lattice² in both conventional and unconventional superconductors. In a previous paper³ (referred to as I) we gave a group theoretical classification of vortex lattice solutions of the BG equation in a uniform magnetic field of the extended Hubbard model on a two-dimensional square lattice. In that paper, we considered conventional definitions^{4,5,6} of s -wave, d -wave and p -wave local symmetric order parameters (OPs), which are valid definitions of OPs in a spatially uniform superconducting state in the absence of a magnetic field. We showed that, e.g., s -wave OP at the site obtained under translation by a lattice vector (of the vortex lattice) from a site (m, n) is always a linear combination of s -wave, d -wave and p -wave OPs at the site (m, n) . We then pointed out that it is not appropriate to assign conventional order parameters at each site.

In this paper we give more appropriate definitions of OPs such that an OP at the translated site is expressed by the corresponding OP at the original site times a phase factor. The model and notation used here are the same as those in I.

We consider singlet superconductivity on a square lattice with inter-atomic spacing a . The Hamiltonian for the system is given by

$$\begin{aligned}
 H = & -t \sum_{(m,n)s} \{ (e^{iKn} a_{(m,n)s}^\dagger a_{(m+1,n)s} + e^{-iKm} a_{(m,n)s}^\dagger a_{(m,n+1)s}) + \text{H.c} \} \\
 & -\mu \sum_{(m,n)s} a_{(m,n)s}^\dagger a_{(m,n)s} \\
 & +U \sum_{(m,n)} a_{(m,n)\uparrow}^\dagger a_{(m,n)\uparrow} a_{(m,n)\downarrow}^\dagger a_{(m,n)\downarrow} \\
 & +V \sum_{(m,n)ss'} \{ a_{(m,n)s}^\dagger a_{(m,n)s} a_{(m+1,n)s'}^\dagger a_{(m+1,n)s'} \\
 & + a_{(m,n)s}^\dagger a_{(m,n)s} a_{(m,n+1)s'}^\dagger a_{(m,n+1)s'} \},
 \end{aligned} \tag{1}$$

where (m, n) denotes a site in a square crystal lattice, $\mathbf{B} = (0, 0, B)$ is a magnetic field, $K = (\pi\phi)/(2\phi_0)$, $\phi = Ba^2$ is the magnetic flux through a unit cell of the crystal lattice, and $\phi_0 = ch/2e$ is the flux quantum. The symmetry group of the system is given by

$$G_0 = (e + tC_{2x})C_4\mathbf{T}\mathbf{S}\Phi, \tag{2}$$

where $\mathbf{T}$ is the group of the magnetic translation,⁷ consisting of the elements $T(Ma\mathbf{e}_x + Na\mathbf{e}_y)$ ($M, N = \text{integer}$), such that

$$T(Mae_x + Nae_y) \cdot a_{(m,n)s}^\dagger = e^{iK(Mn-Nm)} a_{(m+M,n+N)s}^\dagger \quad (3)$$

and $C_4 = (e, C_{4z}^+, C_{2z}, C_{4z}^-)$ is the four-fold rotation group around the origin $(0,0)$. Its generator's action on the Fermion operator is given by

$$C_{4z}^+ \cdot a_{(m,n)s}^\dagger = a_{(-n,m)s}^\dagger. \quad (4)$$

The action of tC_{2x} (t is time reversal)³ on the Fermion operator is

$$\begin{aligned} tC_{2x} \cdot (fa_{(m,n)\uparrow}^\dagger) &= -f^* a_{(m,-n)\downarrow}^\dagger, \\ tC_{2x} \cdot (fa_{(m,n)\downarrow}^\dagger) &= f^* a_{(m,-n)\uparrow}^\dagger, \end{aligned} \quad (5)$$

where f is a complex number. Also in Eq. (2), $\mathbf{S}$ is the group of the spin rotation ($SU(2)$), and Φ is the group of the global gauge transformation.

Hereafter, for illustrative purposes, we restrict our consideration to the symmorphic tetragonal vortex lattice states for the case $\phi = \phi_0/p^2$ (p : integer), in which the center point of each rotation is at a site of the crystal lattice. In this case there are four types of tetragonal vortex lattice solutions of the BG equation characterized by invariance groups³:

$$G_{(l)} = (e + tC_{2x}) \tilde{C}^l \mathbf{L} \mathbf{S} \quad (l = 0, 2, \pm 1). \quad (6)$$

Here $\mathbf{L}$ is a subgroup of $\mathbf{T}\Phi$ consisting elements $L(Mpe_x + Npe_y)$ such that

$$L(Mpe_x + Npe_y) a_{(m,n)s}^\dagger \equiv e^{i\frac{\pi}{2}(MN+M+N)} T(Mpe_x + Npe_y) a_{(m,n)s}^\dagger, \quad (7)$$

and

$$\tilde{C}^l = \sum_j e^{i\frac{2\pi}{4}lj} C_{4z}^j, \quad (8)$$

where the phase factor $e^{i\frac{2\pi}{4}lj}$ should be understood to appear when a group element acts on the space $W(a^\dagger a^\dagger)$. (The case $l = -1$ is overlooked in I.) From $\mathbf{L}$ and $\tilde{C}^l$ invariance of the generalized density matrix we have

$$\begin{aligned} &\langle (L(Mpe_x + Npe_y) a_{(m,n)s}) (L(Mpe_x + Npe_y) a_{(m',n')s'}) \rangle \\ &= \langle a_{(m,n)s} a_{(m',n')s'} \rangle \\ &\langle e^{-i\frac{2\pi}{4}l} (C_{4z}^+ a_{(m,n)s}) (C_{4z}^+ a_{(m',n')s'}) \rangle = \langle a_{(m,n)s} a_{(m',n')s'} \rangle \end{aligned} \quad (9)$$

Then we have

$$\begin{aligned} &\langle a_{(m+Mp,n+Np)s} a_{(m'+Mp,n'+Np)s'} \rangle \\ &= e^{i\pi(MN+M+N)+i\frac{\pi}{2p}(M(n+n')-N(m+m'))} \langle a_{(m,n)s} a_{(m',n')s'} \rangle, \\ &\langle a_{(-n,m)s} a_{(-n',m')s'} \rangle = e^{i\frac{\pi}{2}l} \langle a_{(m,n)s} a_{(m',n')s'} \rangle. \end{aligned} \quad (10)$$

For the case $l = -1, p = 3$, using (10) we obtain a schematic pattern of bond OPs such as $\langle a_{(m,n)\downarrow} a_{(m+1,n)\uparrow} \rangle$, which is displayed in Fig. 1.




FIG. 1. Schematic pattern of bond OP $\langle a_{(m,n)\downarrow} a_{(m+1,n)\uparrow} \rangle$ and $\langle a_{(m,n)\downarrow} a_{(m,n+1)\uparrow} \rangle$ for $p = 3$, $l = -1$. The angle of an arrow with respect to the horizontal x axis represents the phase of the OP. The inner square represents a magnetic unit cell.

It is known^{4,8} that even in the case of an isolated vortex line, a d -wave OP induces an s -wave OP. In I, we use conventional s -wave, d -wave and p -wave OPs^{4,5,6} on a site (m, n) as follows:

$$\begin{aligned}
S(m, n) &= \frac{1}{4} \{ \langle a_{(m,n)\downarrow} a_{(m+1,n)\uparrow} \rangle + \langle a_{(m,n)\downarrow} a_{(m-1,n)\uparrow} \rangle \\
&\quad + \langle a_{(m,n)\downarrow} a_{(m,n+1)\uparrow} \rangle + \langle a_{(m,n)\downarrow} a_{(m,n-1)\uparrow} \rangle \}, \\
D(m, n) &= \frac{1}{4} \{ \langle a_{(m,n)\downarrow} a_{(m+1,n)\uparrow} \rangle + \langle a_{(m,n)\downarrow} a_{(m-1,n)\uparrow} \rangle \\
&\quad - \langle a_{(m,n)\downarrow} a_{(m,n+1)\uparrow} \rangle - \langle a_{(m,n)\downarrow} a_{(m,n-1)\uparrow} \rangle \}, \\
P_x(m, n) &= \frac{1}{2} \{ \langle a_{(m,n)\downarrow} a_{(m+1,n)\uparrow} \rangle - \langle a_{(m,n)\downarrow} a_{(m-1,n)\uparrow} \rangle \}, \\
P_y(m, n) &= \frac{1}{2} \{ \langle a_{(m,n)\downarrow} a_{(m,n+1)\uparrow} \rangle - \langle a_{(m,n)\downarrow} a_{(m,n-1)\uparrow} \rangle \}.
\end{aligned} \tag{11}$$

Then we have shown that these OPs at the site $(m + Mp, n + Np)$ are expressed by a linear combination of s -wave, d -wave and p -wave OPs at the site (m, n) :

$$\begin{aligned}
D(m + Mp, n + Np) &= \frac{1}{2} e^{i\pi(MN+M+N)} e^{i\frac{\pi}{p}(Mn-Nm)} \\
&\quad \times \{ (\cos(\frac{\pi N}{2p}) - \cos(\frac{\pi M}{2p})) S(m, n) \\
&\quad + (\cos(\frac{\pi N}{2p}) + \cos(\frac{\pi M}{2p})) D(m, n) \\
&\quad + i \sin(\frac{\pi N}{2p}) P_x(m, n) - i \sin(\frac{\pi M}{2p}) P_y(m, n) \},
\end{aligned} \tag{12}$$

and so on. Then it is very difficult to recognize the symmetry of magnetic translation of the vortex lattice by inspecting d wave OPs $D(m, n)$ and so on. Thus it is not appropriate to describe a vortex lattice state by assigning the conventional d -wave OP at each site.

Here we give more refined definitions of s -wave, d -wave and p -wave OPs on a site (m, n) . As shown in I, $2\pi/4$, π and $(-2\pi)/4$ rotation around (m, n) (we denote them by $C_{4z}^+(m, n)$, $C_{2z}(m, n)$ and $C_{4z}^-(m, n)$) are given by

$$\begin{aligned}
C_{4z}^+(m, n) &= T(mae_x + nae_y) C_{4z}^+ T(-mae_x - nae_y), \\
C_{2z}(m, n) &= T(mae_x + nae_y) C_{2z} T(-mae_x - nae_y), \\
C_{4z}^-(m, n) &= T(mae_x + nae_y) C_{4z}^- T(-mae_x - nae_y)
\end{aligned} \tag{13}$$

Thus we have

$$\begin{aligned}
C_{4z}^+(m, n) \cdot a_{(m+1, n)s} &= e^{-iK(m+n)} a_{(m, n+1)s}, \\
C_{2z}(m, n) \cdot a_{(m+1, n)s} &= e^{-2iKn} a_{(m-1, n)s}, \\
C_{4z}^-(m, n) \cdot a_{(m+1, n)s} &= e^{-iK(n-m)} a_{(m, n-1)s}, \\
C_{4z}^+(m, n) \cdot a_{(m, n)s} &= C_{2z}(m, n) \cdot a_{(m, n)s} = C_{4z}^-(m, n) \cdot a_{(m, n)s} \\
&= a_{(m, n)s}.
\end{aligned} \tag{14}$$

Here we define s -wave, d -wave and p -wave OPs on site (m, n) as A, B and E representation of the four fold rotation group $C_4(m, n)$ around the point (m, n) consisting of $\{e, C_{4z}^+(m, n), C_{2z}(m, n)$ and $C_{4z}^-(m, n)\}$ as follows :

$$\begin{aligned}
\tilde{S}(m, n) &= \frac{1}{4} e^{iKn} \{ \langle a_{(m, n)\downarrow} a_{(m+1, n)\uparrow} \rangle + \langle (C_{4z}^+(m, n) a_{(m, n)\downarrow}) (C_{4z}^+(m, n) a_{(m+1, n)\uparrow}) \rangle \\
&\quad + \langle (C_{2z}(m, n) a_{(m, n)\downarrow}) (C_{2z}(m, n) a_{(m+1, n)\uparrow}) \rangle \\
&\quad + \langle (C_{4z}^-(m, n) a_{(m, n)\downarrow}) (C_{4z}^-(m, n) a_{(m+1, n)\uparrow}) \rangle \}, \\
\tilde{D}(m, n) &= \frac{1}{4} e^{iKn} \{ \langle a_{(m, n)\downarrow} a_{(m+1, n)\uparrow} \rangle - \langle (C_{4z}^+(m, n) a_{(m, n)\downarrow}) (C_{4z}^+(m, n) a_{(m+1, n)\uparrow}) \rangle \\
&\quad + \langle (C_{2z}(m, n) a_{(m, n)\downarrow}) (C_{2z}(m, n) a_{(m+1, n)\uparrow}) \rangle \\
&\quad - \langle (C_{4z}^-(m, n) a_{(m, n)\downarrow}) (C_{4z}^-(m, n) a_{(m+1, n)\uparrow}) \rangle \}, \\
\tilde{P}_x(m, n) &= \frac{1}{2} e^{iKn} \{ \langle a_{(m, n)\downarrow} a_{(m+1, n)\uparrow} \rangle - \langle (C_{2z}(m, n) a_{(m, n)\downarrow}) (C_{2z}(m, n) a_{(m+1, n)\uparrow}) \rangle \}, \\
\tilde{P}_y(m, n) &= \frac{1}{2} e^{iKn} \{ \langle (C_{4z}^+(m, n) a_{(m, n)\downarrow}) (C_{4z}^+(m, n) a_{(m+1, n)\uparrow}) \rangle \\
&\quad - \langle (C_{4z}^-(m, n) a_{(m, n)\downarrow}) (C_{4z}^-(m, n) a_{(m+1, n)\uparrow}) \rangle \}.
\end{aligned} \tag{15}$$

Here the factor e^{iKn} is used to obtain a symmetrical expression. Using (14) we have

$$\begin{aligned}
\tilde{S}(m, n) &= \frac{1}{4} \{ e^{iKn} \langle a_{(m, n)\downarrow} a_{(m+1, n)\uparrow} \rangle + e^{-iKm} \langle a_{(m, n)\downarrow} a_{(m, n+1)\uparrow} \rangle \\
&\quad + e^{-iKn} \langle a_{(m, n)\downarrow} a_{(m-1, n)\uparrow} \rangle + e^{iKm} \langle a_{(m, n)\downarrow} a_{(m, n-1)\uparrow} \rangle \}, \\
\tilde{D}(m, n) &= \frac{1}{4} \{ e^{iKn} \langle a_{(m, n)\downarrow} a_{(m+1, n)\uparrow} \rangle - e^{-iKm} \langle a_{(m, n)\downarrow} a_{(m, n+1)\uparrow} \rangle \\
&\quad + e^{-iKn} \langle a_{(m, n)\downarrow} a_{(m-1, n)\uparrow} \rangle - e^{iKm} \langle a_{(m, n)\downarrow} a_{(m, n-1)\uparrow} \rangle \}, \\
\tilde{P}_x(m, n) &= \frac{1}{2} \{ e^{iKn} \langle a_{(m, n)\downarrow} a_{(m+1, n)\uparrow} \rangle - e^{-iKn} \langle a_{(m, n)\downarrow} a_{(m-1, n)\uparrow} \rangle \}, \\
\tilde{P}_y(m, n) &= \frac{1}{2} \{ e^{-iKm} \langle a_{(m, n)\downarrow} a_{(m, n+1)\uparrow} \rangle \\
&\quad - e^{iKm} \langle a_{(m, n)\downarrow} a_{(m, n-1)\uparrow} \rangle \}
\end{aligned} \tag{16}$$

From (10) we can see that

$$\begin{aligned}
\tilde{S}(m + Mp, n + Np) &= e^{i\pi(MN+M+N)} e^{i\frac{\pi}{p}(Mn-Nm)} \tilde{S}(m, n), \\
\tilde{D}(m + Mp, n + Np) &= e^{i\pi(MN+M+N)} e^{i\frac{\pi}{p}(Mn-Nm)} \tilde{D}(m, n), \\
\tilde{P}_x(m + Mp, n + Np) &= e^{i\pi(MN+M+N)} e^{i\frac{\pi}{p}(Mn-Nm)} \tilde{P}_x(m, n), \\
\tilde{P}_y(m + Mp, n + Np) &= e^{i\pi(MN+M+N)} e^{i\frac{\pi}{p}(Mn-Nm)} \tilde{P}_y(m, n) \\
\tilde{S}(-n, m) &= e^{i\frac{\pi}{2}l} \tilde{S}(m, n), \\
\tilde{D}(-n, m) &= -e^{i\frac{\pi}{2}l} \tilde{D}(m, n), \\
\tilde{P}_x(-n, m) &= -e^{i\frac{\pi}{2}l} \tilde{P}_y(m, n), \\
\tilde{P}_y(-n, m) &= e^{i\frac{\pi}{2}l} \tilde{P}_x(m, n)
\end{aligned} \tag{17}$$

Thus these new OPs do not mix due to the invariance magnetic translation $\mathbf{L}$. In Figs. 2 and 3 we show schematic patterns of $\tilde{S}(m, n)$ and $\tilde{D}(m, n)$ corresponding to Fig. 1 for the case $l = -1, p = 3$.




FIG. 2. A schematic pattern of $\tilde{S}(m, n)$ for the case $p = 3, l = -1$.




FIG. 3. A schematic pattern of $\tilde{D}(m, n)$ for the case $p = 3, l = -1$.

In the process of making one rotation around the origin $(0, 0)$: $(m, n) \Rightarrow (-n, m) \Rightarrow (-m, -n) \Rightarrow (n, -m) \Rightarrow (n, m)$, total change of the phase of the s -wave OP is $2\pi l$. The corresponding total change

of phase for the d -wave OP is $2\pi(l \pm 2)$. Thus we obtain winding numbers of s -wave and d -wave OPs in Table I. Thus we see that s -wave and d -wave have opposite winding numbers for the cases $l = +1$ and $l = -1$. The opposite winding of s -wave and d -wave OPs has been pointed by Volovik⁸ in an isolated vortex, which corresponds to the case $l = -1$. Note that this relation holds also in the cases of vortex lattice for $l = \pm 1$.

TABLE I. Winding number N_W around $(0, 0)$.

l	$N_W(\tilde{S})$	$N_W(\tilde{D})$
0	0	2
2	2	0
+1	+1	-1
-1	-1	+1

In conclusion, we obtained well-defined OPs at a site (m, n) : $\tilde{S}(m, n)$, $\tilde{D}(m, n)$, $\tilde{P}_x(m, n)$ and $\tilde{P}_y(m, n)$, which have a well-defined nature such that an OP at the site obtained under translation by a basis vector (of the vortex lattice) from a site (m, n) is expressed by the corresponding OP at the site (m, n) times a phase factor. Similar group theoretical analysis also works for the case in which the center of the vortex is located in the middle of a plaquette of the crystal lattice.

The authors would like to express their thanks to M. Ichioka for helpful discussions.

* e-mail: ozaki@cc.kochi-u.ac.jp

† e-mail: hori@ge.kochi-ct.ac.jp

‡ e-mail: goto@ge.kochi-ct.ac.jp

¹ N. Hayashi, M. Ichioka and K. Machida, J. Phys. Soc. Jpn. **67** (1998), 3367.

² T. Kita, J. Phys. Soc. Jpn. **67** (1998), 2075.

³ M. Ozaki, M. Yamazaki, A. Goto and Y. Hori, Prog. Theor. Phys. **100** (1998), 253.

⁴ P. I. Soininen, C. Kallin and A. J. Berlinsky, Phys. Rev. **B50** (1994), 13883.

⁵ Y. Wang and A. H. MacDonald, Phys. Rev. **B52** 1995, R3876.

⁶ H. Tanaka, K. Kuboki and M. Sigrist, Int. J. Mod. Phys. **12** (1998), 2447.

⁷ E. Brown, Phys. Rev. **133** (1964), A1038.

⁸ G. E. Volovik, JETP Lett. **58** (1993), 469.